

SEMIGROUPS IN WHICH THE RADICAL OF EVERY IDEAL IS A SUBSEMIGROUP

STOJAN BOGDANOVIĆ AND MIROSLAV ĆIRIĆ

Abstract. *In this paper we consider semigroups S in which $\sqrt{A} = \{x \in S \mid (\exists n \in \mathbf{Z}^+) x^n \in A\}$ is a subsemigroup of S for every ideal A of S .*

1. Introduction and preliminaries

Throughout paper, by \mathbf{Z}^+ we denote the set of all positive integers. If $a, b \in S$, then $a \mid b$ iff $b = xay$ for some $x, y \in S^1$, $a \overset{r}{\mid} b$ iff $ax = b$ for some $x \in S^1$, $a \underset{l}{\mid} b$ iff $xa = b$ for some $x \in S^1$, $a \overset{r}{\mid} b$ iff $a \overset{r}{\mid} b$ and $a \underset{l}{\mid} b$, $a \longrightarrow b$ iff $a \mid b^i$ for some $i \in \mathbf{Z}^+$ and $a \overset{h}{\longrightarrow} b$ iff $a \overset{h}{\mid} b^i$ for some $i \in \mathbf{Z}^+$, where h is r, l or t . A semigroup S is *Archimedean* (*right Archimedean*, *t-Archimedean*, *power joined*) iff for all $a, b \in S$, $a \longrightarrow b$ ($a \overset{r}{\longrightarrow} b$, $a \overset{t}{\longrightarrow} b$, $\langle a \rangle \cap \langle b \rangle \neq \emptyset$). By the *radical* of the subset A of a semigroup S we mean the set \sqrt{A} defined by

$$\sqrt{A} = \{x \in S \mid (\exists n \in \mathbf{Z}^+) x^n \in A\}.$$

If S is a semigroup with the zero 0 , then an element $a \in S$ is a *nilpotent* if there exists $n \in \mathbf{Z}^+$ such that $a^n = 0$, and the set of all nilpotents of S we denote by $Nil(S)$.

T.Tamura and N.Kimura [12] showed that every commutative semigroup is a semilattice of Archimedean semigroups. This well known result has since been generalized by many authors [2-13]. Semigroups which are semilattice of Archimedean semigroups are completely described by M.S.Putcha [7], T.Tamura [10] and by M.Ćirić and S.Bogdanović [4]. M.S.Putcha [7] has proved the following

THEOREM P. *A semigroup S is a semilattice of Archimedean semigroups if and only if*

$$a \mid b \Rightarrow a^2 \longrightarrow b$$

for all $a, b \in S$. \square

These semigroups are, also, completely described by M. Ćirić and S. Bogdanović in [4] by the following

THEOREM ĆB. *The following conditions are equivalent on a semigroup S :*

- (i) S is a semilattice of Archimedean semigroups;
- (ii) $(\forall a, b \in S) a^2 \rightarrow ab$;
- (iii) the radical of every ideal of S is an ideal of S . \square

Л.Н.Шеврин [14] showed that the equivalence (i) \Leftrightarrow (iii) of Theorem ĆB. holds if S is completely π -regular $((\forall a \in S)(\exists x \in S)(\exists n \in \mathbf{Z}^+) a^n = a^n x a^n, a^n x = x a^n)$.

In this paper we characterize semigroups in which the radical of every ideal (right ideal, bi-ideal, subsemigroup) is a subsemigroup (or ideal or bi-ideal or right ideal).

For undefined notion and notations we refer to [1].

2. Main results

THEOREM 1. *The following conditions on a semigroup S are equivalent:*

- (i) the radical of every ideal of S is a subsemigroup of S ;
- (ii) in every homomorphic image with zero of S the set of all nilpotent elements is a subsemigroup;
- (iii) $(\forall a, b \in S)(\forall k, l \in \mathbf{Z}^+) a^k \rightarrow ab \vee b^l \rightarrow ab$.

PROOF. (i) \Rightarrow (iii). Let $a, b \in S, k, l \in \mathbf{Z}^+$. Since $A = S\{a^k, b^l\}S$ is an ideal of S and $a, b \in \sqrt{A}$ we then have by hypothesis that $ab \in \sqrt{A}$. Hence, there exists $n \in \mathbf{Z}^+$ such that $(ab)^n \in S\{a^k, b^l\}S$. Thus $a^k \rightarrow ab$ or $b^l \rightarrow ab$.

(iii) \Rightarrow (ii). Let T be a semigroup with the zero element and let T be a homomorphic image of S . Then the condition (iii) holds in T . For every $a, b \in Nil(T)$ there exist $m, l \in \mathbf{Z}^+$ such that $a^m = b^l = 0$, and thus $(ab)^n \in T\{0, 0\}T = \{0\}$, for some $n \in \mathbf{Z}^+$. Therefore, $Nil(T)$ is a subsemigroup of T .

(ii) \Rightarrow (i). Let A be an ideal of S . Let φ be a homomorphism of S onto S/A . Let $a, b \in \sqrt{A}$. Since $\varphi(a), \varphi(b) \in Nil(S/A)$ we then have that $\varphi(a)\varphi(b) \in Nil(S/A)$, i.e. $\varphi(ab) \in Nil(S/A)$ and thus $(ab)^n \in A$ for some $n \in \mathbf{Z}^+$. Hence $ab \in \sqrt{A}$, i.e. \sqrt{A} is a subsemigroup of S . \square

In a similar way as in the previous theorem it can be proved the following

THEOREM 2. \sqrt{R} is a subsemigroup of S , for every right ideal R of S if and only if

$$(\forall a, b \in S)(\forall k, l \in \mathbf{Z}^+) a^k \xrightarrow{r} ab \vee b^l \xrightarrow{r} ab. \square$$

THEOREM 3. *The following conditions on a semigroup S are equivalent:*

- (i) S is a semilattice of Archimedean semigroups;
- (ii) $(\forall a, b \in S) a \underset{r}{|} b \Rightarrow a^2 \rightarrow b$;

- (iii) \sqrt{SaS} is an ideal of S , for all $a \in S$;
- (iv) in every homomorphic image with zero of S the set of all nilpotent elements is an ideal.

PROOF. By Theorems **P.** and **ĆB.** \square

THEOREM 4. *The radical of every right ideal of a semigroup S is a bi-ideal of S if and only if*

$$(1) \quad (\forall a, b, c \in S)(\forall k, l \in \mathbf{Z}^+) a^k \xrightarrow{r} abc \quad \vee \quad c^l \xrightarrow{r} abc .$$

PROOF. Let $a, b, c \in S$ and let $k, l \in \mathbf{Z}^+$. Assume that $R = \{a^k, c^l\}S$. Since $a, c \in \sqrt{R}$ and \sqrt{R} is a bi-ideal of S we then have that $abc \in \sqrt{R}S\sqrt{R} \subseteq \sqrt{R}$, i.e. there exists $n \in \mathbf{Z}^+$ such that $(abc)^n \in R = \{a^k, c^l\}S$. Thus $a^k \xrightarrow{r} abc$ or $c^l \xrightarrow{r} abc$.

Conversely, let R be a right ideal of S . For $a, c \in \sqrt{R}$ there exist $k, l \in \mathbf{Z}^+$ such that $a^k, c^l \in R$. Now by (1) we have that

$$\{a^k, c^l\}S \subseteq RS \subseteq R$$

for some $n \in \mathbf{Z}^+$. Hence, $abc \in \sqrt{R}$. Therefore, \sqrt{R} is a bi-ideal of S . \square

THEOREM 5. *The following conditions on a semigroup S are equivalent:*

- (i) S is a semilattice of right Archimedean semigroups;
- (ii) $(\forall a, b \in S)(\forall k \in \mathbf{Z}^+) b^k \xrightarrow{r} ab$;
- (iii) the radical of every right ideal of S is a left ideal of S .

PROOF. (i) \Rightarrow (ii). Let S be a semilattice Y of right Archimedean semigroups S_α , $\alpha \in Y$. Then for $a \in S_\alpha$, $b \in S_\beta$ we have that $ab, b^k a \in S_{\alpha\beta}$, for all $k \in \mathbf{Z}^+$, and there exists $n \in \mathbf{Z}^+$ such that

$$(ab)^n \in b^k a S_{\alpha\beta} \subseteq b^k S .$$

Thus $b^k \xrightarrow{r} ab$.

(ii) \Rightarrow (i). This implication follows by Proposition 1.1.[2].

(ii) \Rightarrow (iii). Let R be a right ideal of S . Assume that $a \in S$, $b \in \sqrt{R}$. Then $b^k \in R$, for some $k \in \mathbf{Z}^+$, and we have that

$$(ab)^n \in b^k S \subseteq RS \subseteq R ,$$

for some $n \in \mathbf{Z}^+$. Thus $ab \in \sqrt{R}$, i.e. \sqrt{R} is a left ideal of S .

(iii) \Rightarrow (i). Let $a, b \in S$, $R = bS$. Then $b \in \sqrt{R}$. Since \sqrt{R} is a left ideal of S we then have that $ab \in \sqrt{R}$, i.e. there exists $n \in \mathbf{Z}^+$ such that $(ab)^n \in R = bS$, whence by Proposition 1.1.[2] we have that the condition (i) holds. \square

THEOREM 6. *The following conditions on a semigroup S are equivalent:*

- (i) $(\forall a, b \in S) a \underset{r}{|} b \Rightarrow a^2 \xrightarrow{r} b$;
- (ii) $(\forall a, b \in S)(\forall k \in \mathbf{Z}^+) a^k \xrightarrow{r} ab$;
- (iii) $(\forall a, b \in S) a^2 \xrightarrow{r} ab$;

- (iv) \sqrt{aS} is a right ideal of S , for every $a \in S$;
 (v) \sqrt{R} is a right ideal of S , for every right ideal R of S .

PROOF. (i) \Rightarrow (iii). Since $ab \in aS$ for every $a, b \in S$, we then have that $(ab)^n \in a^2S$. Thus $a^2 \xrightarrow{r} ab$.

(iii) \Rightarrow (ii). By induction.

(ii) \Rightarrow (i). Let $b = au$ for some $u \in S$. Then there exists $n \in \mathbf{Z}^+$ such that $b^n = (au)^n \in a^2S$. Thus $a^2 \xrightarrow{r} b$.

(ii) \Rightarrow (iv). Let $x \in \sqrt{aS}$ and let $b \in S$. Then $x^k \in aS$ for some $k \in \mathbf{Z}^+$. Since

$$(xb)^n \in x^k S \subseteq aSS \subseteq aS, \text{ for some } n \in \mathbf{Z}^+$$

we then have that $xb \in \sqrt{aS}$. Thus \sqrt{aS} is a right ideal of S .

(iv) \Rightarrow (iii). Let $a, b \in S$. Then $a \in \sqrt{a^2S}$. Since $\sqrt{a^2S}$ is a right ideal of S , then $ab \in \sqrt{a^2S}$, and therefore (iii) holds.

(v) \Rightarrow (iv). Since aS is a right ideal of S , by (v) we then have that \sqrt{aS} is also right ideal of S .

(ii) \Rightarrow (v). Let R be a right ideal of S . Let $a \in \sqrt{R}$, $b \in S$. Then $a^k \in R$ for some $k \in \mathbf{Z}^+$. Now,

$$(ab)^n \in a^k S \subseteq RS \subseteq R \text{ for some } n \in \mathbf{Z}^+$$

and thus $ab \in \sqrt{R}$, i.e. \sqrt{R} is a right ideal of S . \square

THEOREM 7. The following conditions on a semigroup S are equivalent:

- (i) S is a normal band of t -Archimedean semigroups;
 (ii) $(\forall a, b, c \in S) ac \xrightarrow{t} abc$;
 (iii) for every $a, b, c \in S$,

$$a \underset{r}{|} c \wedge b \underset{l}{|} c \Rightarrow ab \xrightarrow{t} c.$$

PROOF. (i) \Leftrightarrow (ii). This equivalence is from [3].

(ii) \Rightarrow (iii). Let $a \underset{r}{|} c$, $b \underset{l}{|} c$. then there exist $u, v \in S$ such that $c = au = vb$, whence $c^2 = auvb$. Now, there exists $n \in \mathbf{Z}^+$ such that $c^{2n} = (auvb)^n \in abSab$, i.e. $ab \xrightarrow{t} c$.

(iii) \Rightarrow (ii). It is clear that $a \underset{r}{|} abc$, $c \underset{l}{|} abc$, for every $a, b, c \in S$. By (iii) there exists $n \in \mathbf{Z}^+$ such that $(abc)^n \in acS \cap Sac$. Hence, (ii) holds. \square

THEOREM 8. The following conditions on a semigroup S are equivalent:

- (i) $(\forall a, b \in S) a^2b \xrightarrow{r} ab$;
 (ii) for all $a, b, c \in S$,

$$a \underset{r}{|} c \wedge b \underset{r}{|} c \Rightarrow ab \xrightarrow{r} c.$$

PROOF. (i) \Rightarrow (ii). Let $c = au = bv$ for some $u, v \in S$, whence $c^2 = (au)^2$. Now, there exists $i \in \mathbf{Z}^+$ such that

$$c^{2i} = (a(au))^i \in a^2uauS \subseteq a^2uS = a(au)S = a(bv)S \subseteq abS.$$

Thus $ab \xrightarrow{r} c$.

(ii) \Rightarrow (i). It is clear that $a \underset{r}{|} ab, ab \underset{r}{|} ab$, for all $a, b \in S$, and by (ii) we have that $a(ab) = a^2b \xrightarrow{r} ab$. \square

THEOREM 9. *The following conditions on a semigroup S are equivalent:*

- (i) S is a semilattice of t -Archimedean semigroups;
- (ii) $(\forall a, b \in S)(\exists n \in \mathbf{Z}^+) (ab)^n \in bSa$;
- (iii) the radical of every bi-ideal of S is an ideal of S .

PROOF. (i) \Leftrightarrow (ii). This equivalence is from [2].

(ii) \Rightarrow (iii). Let A be a bi-ideal of S and let $a \in \sqrt{A}$, $b \in S$. Then $a^k \in A$ for some $k \in \mathbf{Z}^+$, whence

$$(ab)^m, (ba)^n \in a^k b S b a^k \subseteq A S A \subseteq A,$$

for some $m, n \in \mathbf{Z}^+$. Thus $ab, ba \in \sqrt{A}$, i.e. \sqrt{A} is an ideal of S .

(iii) \Rightarrow (ii). Let $a, b \in S$. Assume that $A = aSa$, $B = bSb$. It is clear that $a \in \sqrt{A}$, $b \in \sqrt{B}$. Since \sqrt{A} and \sqrt{B} are ideals of S , we then have that $ab \in \sqrt{A} \cap \sqrt{B}$, i.e. there exist $m, n \in \mathbf{Z}^+$ such that $(ab)^m \in aSa$, $(ab)^n \in bSb$, whence $(ab)^{m+n} \in bSbaSa \subseteq bSa$. \square

3. More on bi-ideals and radicals

The Theorems 10.-22. can be proved as some of the previous theorems and the proof of any of its will be omitted.

THEOREM 10. *The radical of every ideal of a semigroup S is a bi-ideal of S if and only if*

$$(\forall a, b, c \in S)(\forall k, l \in \mathbf{Z}^+) a^k \longrightarrow abc \quad \vee \quad c^l \longrightarrow abc. \quad \square$$

THEOREM 11. *The radical of every subsemigroup of a semigroup S is a bi-ideal of S if and only if*

$$(\forall a, b, c \in S)(\forall k, l \in \mathbf{Z}^+)(\exists n \in \mathbf{Z}^+) (abc)^n \in \langle a^k, c^l \rangle. \quad \square$$

THEOREM 12. *The radical of every bi-ideal of a semigroup S is a bi-ideal of S if and only if*

$$(\forall a, b, c \in S)(\forall k, l \in \mathbf{Z}^+)(\exists n \in \mathbf{Z}^+) (abc)^n \in \{a^k, c^l\}S\{a^k, c^l\}. \quad \square$$

THEOREM 13. *The radical of every subsemigroup of a semigroup S is a subsemigroup of S if and only if*

$$(\forall a, b \in S)(\forall k, l \in \mathbf{Z}^+)(\exists n \in \mathbf{Z}^+) (ab)^n \in \langle a^k, b^l \rangle. \quad \square$$

THEOREM 14. *The radical of every bi-ideal of a semigroup S is a subsemigroup of S if and only if*

$$(\forall a, b \in S)(\forall k, l \in \mathbf{Z}^+)(\exists n \in \mathbf{Z}^+) (ab)^n \in \{a^k, b^l\}S\{a^k, b^l\}. \quad \square$$

THEOREM 15. *The radical of every subsemigroup of a semigroup S is a left ideal of S if and only if S is a right zero band of power joined semigroups. \square*

THEOREM 16. *The radical of every subsemigroup of a semigroup S is an ideal of S if and only if S is power joined. \square*

THEOREM 17. *The following conditions on a semigroup S are equivalent:*

- (i) $(\forall a, b, c, d \in S) b^2 \longrightarrow abcd \quad \vee \quad c^2 \longrightarrow abcd;$
(ii) for all $a, b, c \in S,$
 $ab \mid c \Rightarrow a^2 \longrightarrow c \quad \vee \quad b^2 \longrightarrow c. \square$

THEOREM 18. *The following conditions on a semigroup S are equivalent:*

- (i) $(\forall a, b, c \in S) a^2 \xrightarrow{r} abc \quad \vee \quad b^2 \xrightarrow{r} abc;$
(ii) for all $a, b, c \in S,$
 $ab \mid c \Rightarrow a^2 \xrightarrow[r]{} c \quad \vee \quad b^2 \xrightarrow[r]{} c. \square$

THEOREM 19. *The following conditions on a semigroup S are equivalent:*

- (i) $(\forall a, b \in S) a^2 \xrightarrow{t} aba;$
(ii) for all $a, b \in S,$
 $a \mid b \Rightarrow a^2 \xrightarrow[t]{} b. \square$

REFERENCES

- [1] S.BOGDANOVIĆ, *Semigroups with a system of subsemigroups*, Inst. of Math. Novi Sad, 1985.
- [2] S.BOGDANOVIĆ, *Semigroups of Galbiati-Veronesi*, Proc. of the conf. "Algebra and Logic", Zagreb, (1984), 9-20, Novi Sad 1985.
- [3] S.BOGDANOVIĆ AND M.ĆIRIĆ, *Semigroups of Galbiati-Veronesi IV*, Facta Universitatis (Niš), Ser. Math. Inform. (to appear).
- [4] M.ĆIRIĆ AND S.BOGDANOVIĆ, *Decompositions of semigroups induced by identities*, Semigroup Forum (to appear).
- [5] J.L.CHRISLOCK, *On medial semigroups*, J. Algebra **12** (1969), 1-9.
- [6] T.NORDAHL, *Semigroup satisfying $(xy)^m = x^m y^m$* , Semigroup Forum **8** (1974), 332-346.
- [7] M.S.PUTCHA, *Semilattice decomposition of semigroups*, Semigroup Forum, **6** (1973), 12-34.
- [8] M.S.PUTCHA, *Bands of t -Archimedean semigroups*, Semigroup Forum, **6** (1973), 232-239.
- [9] M.S.PUTCHA, *Rings which are semilattices of Archimedean semigroups*, Semigroup Forum, **23** (1981), 1-5.
- [10] T.TAMURA, *On Putcha's theorem concerning semilattice of Archimedean semigroups*, Semigroup Forum, **4** (1972), 83-86.

-
- [11] T.TAMURA, *Quasi-orders, generalized archimedeaness, semilattice decomposition*, Math. Nachr. **68** (1975), 201-220.
- [12] T.TAMURA AND N.KIMURA, *On decomposition of a commutative semigroup*, Kodai Math. Sem. Rep. **4** (1954), 109-112.
- [13] T.TAMURA AND J.SHAFFER, *On exponential semigroups I*, Proc. Japan Acad. **48** (1972), 77-80.
- [14] Л.Н.ШЕВРИН, *Квазипериодические полугруппы, разложимые в связку Архимедовых полугрупп*, XVI Всесоюзн. алгебр. конф. Тезисы докл., Л., 1981, Ч1, с.188.

Faculty of Economics
18000 Niš, Trg Jna 11
Yugoslavia

Department of Mathematics
Philosophical Faculty
18000 Niš, Ћirila i Metodija 2
Yugoslavia

CURRENT ADDRESS: STOJAN BOGDANOVIĆ, FACULTY OF ECONOMICS, 18000 NIŠ, YUGOSLAVIA