

**LACUNARY STRONG CONVERGENCE OF DIFFERENCE
SEQUENCES WITH RESPECT TO A MODULUS
FUNCTION**

RIFAT ÇOLAK

ABSTRACT. A sequence $\theta = (k_r)$ of positive integers is called lacunary if $k_0 = 0$, $0 < k_r < k_{r+1}$ and $h_r = k_r - k_{r-1} \rightarrow \infty$ as $r \rightarrow \infty$. The intervals determined by θ are denoted by $I_r = (k_{r-1}, k_r]$.

Let ω be the set of all sequences of complex numbers and f be a modulus function. Then we define

$$N_\theta(\Delta^m, f) = \{x \in \omega : \lim_r \frac{1}{h_r} \sum_{k \in I_r} f(|\Delta^m x_k - l|) = 0 \text{ for some } l\}.$$

$$N_\theta^0(\Delta^m, f) = \{x \in \omega : \lim_r \frac{1}{h_r} \sum_{k \in I_r} f(|\Delta^m x_k|) = 0\}.$$

$$N_\theta^\infty(\Delta^m, f) = \{x \in \omega : \sup_r \frac{1}{h_r} \sum_{k \in I_r} f(|\Delta^m x_k|) < \infty\}.$$

where $\Delta x_k = x_k - x_{k+1}$, $\Delta^m x_k = \Delta^{m-1} x_k - \Delta^{m-1} x_{k+1}$ and m is a fixed positive integer.

In this study we give various properties and inclusion relations on these sequence spaces.

1. INTRODUCTION

Let w be the set of all sequences of real or complex numbers and ℓ_∞ , c , and c_0 be respectively the Banach spaces of bounded, convergent and null sequences $x = (x_k)$ with the usual norm $\|x\| = \sup_k |x_k|$, where $k \in N = \{1, 2, \dots\}$ the set of positive integers.

The difference sequence space $X(\Delta)$ was introduced by Kizmaz [4] as follows:

$$X(\Delta) = \{x = (x_k) \in w : (\Delta x_k) \in X\} \text{ for } X = \ell_\infty, c \text{ and } c_0,$$

where $\Delta x_k = x_k - x_{k+1}$, for all $k \in N$.

1991 *Mathematics Subject Classification*. Primary: 40C05 , Secondary: 46A45 .
Key words and phrases. Difference sequence, lacunary sequence, modulus function.

The difference sequence spaces were generalized by Et and Çolak [2] as follows:

$$X(\Delta^m) = \{x = (x_k) \in w : \Delta^m x = (\Delta^m x_k) \in X\} \text{ for } X = \ell_\infty, c \text{ and } c_0,$$

where $\Delta^m x_k = \Delta^{m-1} x_k - \Delta^{m-1} x_{k+1}$ and so $\Delta^m x_k = \sum_{v=0}^m (-1)^v \binom{m}{v} x_{k+v}$.

By a lacunary sequence $\theta = (k_r)$; $r = 0, 1, 2, \dots$, where $k_0 = 0$, we mean an increasing sequence of non-negative integers with $h_r = k_r - k_{r-1} \rightarrow \infty$ as $r \rightarrow \infty$. We denote by $I_r = (k_{r-1}, k_r]$ the intervals determined by θ and $q_r = \frac{k_r}{k_{r-1}}$ for $r = 0, 1, 2, \dots$. The space of lacunary convergent sequences N_θ was defined by Freedman et al [3] as follows:

$$(1.1) \quad N_\theta = \left\{ x = (x_k) : \lim_r h_r^{-1} \sum_{k \in I_r} |x_k - \ell| = 0, \text{ for some } \ell \right\}.$$

The space N_θ is a BK-space with the norm

$$(1.2) \quad \|x\|_\theta = \sup_r h_r^{-1} \sum_{k \in I_r} |x_k|.$$

N_θ^0 denotes the subset of N_θ those sequences for which $\ell = 0$ in the definition of N_θ . $(N_\theta^0, \|\cdot\|_\theta)$ is also a BK-space. There is a relation (see for instance [3]) between N_θ and the space $|\sigma_1|$ of strongly Cesaro summable sequences defined by

$$(1.3) \quad |\sigma_1| = \left\{ x = (x_k) : \lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n |x_k - \ell| = 0, \text{ for some } \ell \right\}.$$

In the special case $\theta = (2^r)$ we have $N_\theta = |\sigma_1|$.

The notion of a modulus function was introduced by Nakano [7]. We recall that a modulus f is a function from $[0, \infty)$ to $[0, \infty)$ such that

- (i) $f(x) = 0$ if and only if $x = 0$
- (ii) $f(x + y) \leq f(x) + f(y)$, for all $x \geq 0, y \geq 0$
- (iii) f increasing,
- (iv) f continuous from the right at zero.

Since $|f(x) - f(y)| \leq f(|x - y|)$, it follows from (iv) that f is continuous on $[0, \infty)$. Furthermore, we have $f(nx) \leq nf(x)$ for all $n \in \mathbb{N}$ from condition (ii), and so $f(x) = f\left(nx \frac{1}{n}\right) \leq nf(x/n)$, hence $\frac{1}{n}f(x) \leq f(x/n)$ for all $n \in \mathbb{N}$.

A modulus may be bounded or unbounded. For example, $f(x) = x^p$ for $0 < p \leq 1$ is unbounded, but $f(x) = \frac{x}{x+1}$ is bounded.

Subsequently modulus function have been studied by Ruckle [9], Connor [1], Maddox [5], Pehlivan and Fisher [8], Malkowsky and Savaş [6].

2. MAIN RESULTS

In this section, we extend Δ^m -lacunary strongly convergent sequences which were defined by B.C. Tripathy, M. Et and S. Mahanta [10], and give the relation between Δ^m -lacunary strong convergence and Δ^m -lacunary strong convergence with respect to a modulus.

Definition 2.1. Let f be a modulus function. We define the following sets

$$\begin{aligned} N_\theta^0(\Delta^m, f) &= \left\{ x \in w : \lim_r h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k|) = 0 \right\} \\ N_\theta(\Delta^m, f) &= \left\{ x \in w : \lim_r h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k - \ell|) = 0, \text{ for some } \ell \right\} \\ N_\theta^\infty(\Delta^m, f) &= \left\{ x \in w : \sup_r h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k|) < \infty \right\}. \end{aligned}$$

If we take $f(x) = x$ then we obtain the sequence spaces $N_\theta^0(\Delta^m)$, $N_\theta(\Delta^m)$ and $N_\theta^\infty(\Delta^m)$ from the above sequence spaces.

Theorem 2.2. *The sets $N_\theta^0(\Delta^m, f)$, $N_\theta(\Delta^m, f)$ and $N_\theta^\infty(\Delta^m, f)$ are linear spaces.*

Proof. We consider only $N_\theta(\Delta^m, f)$. Let $x, y \in N_\theta(\Delta^m, f)$ and $\alpha, \gamma \in \mathbb{C}$. Then there exist positive integers K_α and M_γ such that $|\alpha| \leq K_\alpha$ and $|\gamma| \leq M_\gamma$. From the definition of f and Δ^m we have

$$\begin{aligned} &h_r^{-1} \sum_{k \in I_r} f(|\Delta^m(\alpha x_k + \gamma y_k) - (\alpha \ell_1 + \gamma \ell_2)|) \leq \\ &K_\alpha h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k - \ell_1|) + M_\gamma h_r^{-1} \sum_{k \in I_r} f(|\Delta^m y_k - \ell_2|) \rightarrow 0, \quad r \rightarrow \infty. \end{aligned}$$

Thus $N_\theta(\Delta^m, f)$ is a linear space. □

Lemma 2.3. [8] *Let f be a modulus function and let $0 < \delta < 1$. Then for each $x > \delta$ we have $f(x) \leq 2f(1)\delta^{-1}x$.*

Theorem 2.4. *Let f be a modulus function. Then $N_\theta(\Delta^m) \subset N_\theta(\Delta^m, f)$.*

Proof. Let $x \in N_\theta(\Delta^m)$. Then we have

$$(2.1) \quad \tau_r = h_r^{-1} \sum_{k \in I_r} |\Delta^m x_k - \ell| \rightarrow 0 \quad \text{as } r \rightarrow \infty, \text{ for some } \ell.$$

Let $\varepsilon > 0$ and choose δ with $0 < \delta < 1$ such that $f(u) < \varepsilon$ for every u with $0 \leq u \leq \delta$. Then we can write

$$\begin{aligned} & h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k - \ell|) = \\ & h_r^{-1} \sum_{k \in I_r, |\Delta^m x_k - \ell| \leq \delta} f(|\Delta^m x_k - \ell|) + h_r^{-1} \sum_{k \in I_r, |\Delta^m x_k - \ell| > \delta} f(|\Delta^m x_k - \ell|) \leq \\ & \qquad \qquad \qquad h_r^{-1} (h_r \varepsilon) + h_r^{-1} 2f(1)\delta^{-1} h_r \tau_r \end{aligned}$$

from Lemma 2.3. Therefore $x \in N_\theta(\Delta^m, f)$. \square

Theorem 2.5. *Let f be a modulus function, if $\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \beta > 0$, then $N_\theta(\Delta^m, f) = N_\theta(\Delta^m)$.*

Proof. By Theorem 2.4, we need only show that $N_\theta(\Delta^m, f) \subset N_\theta(\Delta^m)$. Let $\beta > 0$ and $x \in N_\theta(\Delta^m, f)$. Since $\beta > 0$, we have $f(t) \geq \beta t$ for all $t \geq 0$. Hence we have

$$h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k - \ell|) \geq h_r^{-1} \sum_{k \in I_r} \beta |\Delta^m x_k - \ell| = \beta h_r^{-1} \sum_{k \in I_r} |\Delta^m x_k - \ell|.$$

Therefore we have $x \in N_\theta(\Delta^m)$. \square

In Theorem 2.5, the condition $\beta > 0$ can not be omitted. For this consider the following example.

Example 2.6. Let $f(x) = \ln(1+x)$. Then $\beta = 0$. Define $\Delta^m x_k$ to be h_r at the $(k_{r-1} + 1)$ -th term in I_r for every $r \geq 1$ and $x_i = 0$ otherwise. Note that $x \notin \ell_\infty(\Delta^m)$. Then we have

$$h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k|) = h_r^{-1} \ln(1+h_r) \rightarrow 0, \quad \text{as } r \rightarrow \infty$$

and so $x \in N_\theta(\Delta^m, f)$, but

$$h_r^{-1} \sum_{k \in I_r} |\Delta^m x_k| = h_r^{-1} h_r \rightarrow 1, \quad \text{as } r \rightarrow \infty,$$

and so $x \notin N_\theta(\Delta^m)$.

Theorem 2.7. *Let $m \geq 1$ be a fixed integer, then*

- (i) $N_\theta^0(\Delta^{m-1}, f) \subset N_\theta^0(\Delta^m, f)$;
- (ii) $N_\theta(\Delta^{m-1}, f) \subset N_\theta(\Delta^m, f)$;
- (iii) $N_\theta^\infty(\Delta^{m-1}, f) \subset N_\theta^\infty(\Delta^m, f)$;

and the inclusions are strict.

Proof. The proof of the inclusions follows from the following inequality

$$h_r^{-1} \sum_{k \in I_r} f(|\Delta^m x_k|) \leq h_r^{-1} \sum_{k \in I_r} f(|\Delta^{m-1} x_k|) + h_r^{-1} \sum_{k \in I_r} f(|\Delta^{m-1} x_{k+1}|).$$

To show the inclusions are strict, let $\theta = (2^r)$ and $x = k^m$. Then $x \in N_\theta^\infty(\Delta^m, f)$, but $x \notin N_\theta^\infty(\Delta^{m-1}, f)$. If $x = (k^m)$, then $\Delta^m x = (-1)^m m!$ and $\Delta^{m-1} x_k = (-1)^{m+1} r! \left(k + \frac{(m-1)}{2}\right)$. \square

Theorem 2.8. *Let $\theta = (k_r)$ be a lacunary sequence. If $1 < \liminf_r q_r \leq \limsup_r q_r < \infty$, then $|\sigma_1|(\Delta^m, f) = N_\theta(\Delta^m, f)$, where*

$$|\sigma_1|(\Delta^m, f) = \left\{ x \in w : \frac{1}{n} \sum_{k=1}^n f(|\Delta^m x_k - \ell|) = 0, \text{ for some } \ell \right\}.$$

Proof. Suppose that $\liminf_r q_r > 1$, then there exists $\delta > 0$ such that $q_r = \frac{k_r}{k_{r-1}} \geq 1 + \delta$ for all $r \geq 1$. Furthermore we have $\frac{k_r}{h_r} \leq \frac{(1+\delta)}{\delta}$ and $\frac{k_{r-1}}{h_r} \leq \frac{1}{\delta}$, for all $r \geq 1$. Then we may write

$$\begin{aligned} (2.2) \quad \frac{1}{h_r} \sum_{i \in I_r} f(|\Delta^m x_i|) &= \frac{1}{h_r} \sum_{i=1}^{k_r} f(|\Delta^m x_i|) - \frac{1}{h_r} \sum_{i=1}^{k_{r-1}} f(|\Delta^m x_i|) \\ &= \frac{k_r}{h_r} \left(k_r^{-1} \sum_{i=1}^{k_r} f(|\Delta^m x_i|) - \frac{k_{r-1}}{h_r} \left(k_{r-1}^{-1} \sum_{i=1}^{k_{r-1}} f(|\Delta^m x_i|) \right) \right). \end{aligned}$$

Now suppose that $\limsup_r q_r < \infty$ and let $\varepsilon > 0$ be given. Then there exists j_0 such that for every $j \geq j_0$

$$A_j = \frac{1}{h_j} \sum_{i \in I_j} [f(|\Delta^m x_i|)] < \varepsilon.$$

We can also choose a number $M > 0$ such that $A_j \leq M$ for all j . If $\limsup_r q_r < \infty$, then there exists a number $\beta > 0$ such that $q_r < \beta$ for every r . Now let n be any integer with $k_{r-1} < n < k_r$. Then

$$\begin{aligned} n^{-1} \sum_{i=1}^n f(|\Delta^m x_i|) &\leq k_{r-1}^{-1} \sum_{i=1}^{k_r} f(|\Delta^m x_i|) \\ &= k_{r-1}^{-1} \left\{ \sum_{i \in I_1} f(|\Delta^m x_i|) + \dots + \sum_{i \in I_r} f(|\Delta^m x_i|) \right\} \\ &= k_{r-1}^{-1} \left\{ \sum_{j=1}^{j_0} \sum_{i \in I_j} f(|\Delta^m x_i|) + \sum_{j=j_0+1}^r \sum_{i \in I_j} f(|\Delta^m x_i|) \right\} \\ &\leq k_{r-1}^{-1} \left\{ \sum_{j=1}^{j_0} \sum_{i \in I_j} f(|\Delta^m x_i|) + \varepsilon (k_r - k_{j_0}) k_{r-1}^{-1} \right\} \\ &= k_{r-1}^{-1} \{h_1 A_1 + h_2 A_2 + \dots + h_{j_0} A_{j_0}\} \\ &\quad + \varepsilon (k_r - k_{j_0}) k_{r-1}^{-1} \end{aligned}$$

$$\begin{aligned} &\leq k_{r-1}^{-1} \left(\sup_{1 \leq i \leq j_0} A_j \right) k_{j_0} + \varepsilon (k_r - k_{j_0}) k_{r-1}^{-1} \\ &< M k_{r-1}^{-1} k_{j_0} + \varepsilon \beta \end{aligned}$$

which yields that $x \in |\sigma_1|(\Delta^m, f)$. □

REFERENCES

- [1] J.S. Connor, On strong matrix summability with respect to a modulus and statistical convergence, *Canad. Math. Bull.* **(32)** (1989), 194–198
- [2] M. Et and R. Çolak, On some generalized difference sequence spaces, *Soochow J. of Math.* **(21)** (1995), 377–386
- [3] A.R. Freedman, J.J. Sember and M. Raphael, Some Cesaro-type summability spaces, *Proc. London Math. Soc.* **(37(3))** (1978), 508–520
- [4] H. Kizmaz, On certain sequence spaces, *Canad. Math. Bull.* **(24)** (1981), 169–176
- [5] I.J. Maddox, Sequence spaces defined by a modulus, *Math. Proc. Camb. Philo. Soc.* **(100)** (1986), 161–166
- [6] E. Malkowsky and E. Savas, Some λ -sequence spaces defined by a modulus, *Archivum Math.* **(36)** (2000), 219–228
- [7] H. Nakano, Concave modulars, *J. Math. Soc. Japan* **(5)** (1953), 29–49
- [8] S. Pehlivan and B. Fisher, Lacunary strong convergence with respect to a sequence of modulus functions, *Comment Math. Univ. Carolin* **(36)** (1995), 69–76
- [9] W.H. Ruckle, FK spaces in which the sequence of coordinate vectors is bounded, *Canad. J. Math.* **(25)** (1973), 1973–1978
- [10] B.C. Tripathy, M. Et and S. Mahanata, On a class of generalized lacunary difference sequences defined by Orlicz function (Submitted).

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, FIRAT UNIVERSITY,
23119 ELAZIG, TÜRKİYE

E-mail address: rcolak@firat.edu.tr